

Original Article



Isolation and Characterization of Oil Obtained from *Leucaena leucocephala* (Lam.) De Wit

Jaya B. Puri | Abdulbasit Shaikh | Umesh Pravin Dhuldhaj*

Department of Biotechnology, School of Life Sciences, Swami Ramanand Teerth Marathwada University, Nanded 431606, India

**Citation** J.B. Puri, A. Shaikh, U.P. Dhuldhaj. **Isolation and Characterization of Oil Obtained from *Leucaena leucocephala* (Lam.) De Wit.** *Eurasian J. Sci. Technol.*, 2024, 4(2), 57-65. <https://doi.org/10.48309/EJST.2024.418888.1100>**Article info:****Received:** 2023-10-01**Accepted:** 2023-11-11**Available Online:** 2023-12-06**ID:** EJST-2310-1100**Checked for Plagiarism:** Yes**Checked Language:** Yes**Keywords:***Leucaena leucocephala*, Medicinal properties, Oil extraction**ABSTRACT**

In the present study, we have collected the leucocephala seeds which are seasonal and having good amount of oil content. The oil seeds can be used as the one of the adulterant for the coffee. The whole plant having importance for human being, it could be as the food or furniture. The immature seeds, pods and flower buds are used as the salad while the gum obtained from this plant can be used for the confectionary, pharmaceuticals, and cosmetics industry. In some of the region, it is considered as the recalcitrant and attracts rodents. The plant is well-known in traditional medicine and seeds are the rich source of several necessary metabolites such as proteins, tannins, carotene, leucenol, and flavonol glycosides and having therapeutic potential like anti-inflammatory and antidiabetic activity. Hence, in our present investigation, we focus for the collection of seeds, and extraction of oil and its characterizations. The dried pods and seeds were collected from Waghi Village, near Nanded, Maharashtra. The obtained extract were further characterized for the confirmation of oil content by measuring lipid content, specific gravity, density, protein, and glucose content, and further analysis with thin layer chromatography. The obtained extract is also subjected to the anti-microbial activity.

Introduction

White Popinac or white lead tree is typically tropical plant species, grows in warm conditions and this plant scientifically known by the name *Leucaena leucocephala* (Lam.) de Wit belongs to Fabaceae family of plants. The plant members of Fabaceae are supposed to nodule forming; hence, also termed as the Leguminosae, the plants of this family can be easily identified by some key markers such as these plants frequently forms symbiotic

relationship with nitrogen fixing bacteria, leaves are compounds, and alternatively arranged, margin is smooth, flower are bisexual, the fruits are legume or loment types. Cultivation of this plants has been done since long time for the health and other necessary benefits. White Popinac is small shrubby in size, supposed to be fast growing with higher photosynthetic rate, responsible for massive biomass production which accounts for around 30 tons dry matter ha⁻¹.yr⁻¹, on average 3-15

*Corresponding Author: Umesh Pravin Dhuldhaj umeshpd12@gmail.com

meter tall and the maximum height reaches to 20 m, numerous branches, young branches covered with smooth, grey to brown, rusty orange-brown vertical fissures on the bark. This plant could be a better solution for the deforestation and lesser plantations [1]. The plant is short-lived (average 20-40 years) evergreen in nature and mature and immature pods can be at the same time on the plant. The plant is resistant to several extreme conditions such as pH (upto 4.1), drought and salinity. The plants have higher resistance to biotic and abiotic stresses, which shows the greater production of secondary metabolites necessary for it [2]. Medicinal plants have a special place in all Indian therapeutic systems like Unani-Tabbi, Ayurved, Siddha, Homeopathy, etc. [3] and White Popinac have versatile medicinal properties and are treated as the miracle tree for food, forage and timber [4]. This plant has potential use in folk medicine to increase the milk production and weight gain in ruminants [4] and has potential as an insect repellent. The major secondary metabolites and necessary phytochemicals obtained from these plants can be used as antidiabetic [5, 6], anticancer, antimetastatic, antibacterial [7], and antihelminthic. The notable medicinal use of this plant is as a contraceptive and abortifacient potential and also helps to reduce stomachache.

This plant species has great potential to sustain in dry conditions and commonly has versatile applications as per the different sites of global locations [8]. It is to be used as a salad or along with other food materials in countries like Indonesia, India, and Thailand. Young plant parts, leaves, and shoots are majorly used to prepare several food dishes while dry seeds were used as an alternative to coffee powder [9], hence also known by the name bush coffee. The countries like Mexico and Central America, where immature seeds and pods are used as food. The gums secreted by this plant are used as an important ingredient in confectionery, cosmetics, and pharmaceuticals.

In some regions, this plant is also supposed to weed and can easily survive and spread in any season, in each season, this plant spreads

thousands of seeds that travel long ways and attract rodents. The invasion of White popinac observed on roadsides, wastelands, cultivated lands, riverbanks, and forest edges which prevent the survival of herbs, shrubs, and trees and this plant influences the particular habitat in patterns of vegetation, plant succession, and community assembly [10].

The seeds are a good source of oil and can be used for the production of biodiesel, and also it is a rich source of lipid, carbohydrate, and protein [11]. The oil obtained from this plant can be used for bio-membrane modeling against xenobiotic compounds [12] and as an anti-corrosive for mild steel and copper. Plants have multifaceted applications like prevention of soil erosion, fuel wood, timber, fodder, green manure, etc. The major amount of oils are coming from the plants in the form of margarine, salad oil, and specially derived products which are necessary components of food processing and preparations. In the present investigation, we have collected the seeds and extracted the oil content from them. The oil obtained from this plant has therapeutic potential. Therefore, we conducted chemical characterization and bioactivity of the obtained oil product.

Materials and Methods

Collection of Sample

The dried pods and seeds were collected from the village Waghi, near Nanded, Maharashtra. The seed sample was further dried in the sunlight for 10 to 12 days. The dried seeds were distributed in two parts, half of the seeds were kept soaked in water for a week and the remaining portion of the seeds were soaked in boiling water for 2-3 hours.

Oil Extraction

The crude extracts obtained in normal water and boiling water [13] were thoroughly mixed with organic solvent such as petroleum ether.

After proper mixing, with the help of separating funnel upper organic layer were separated and subjected to distillation for evaporation of excess solvent present in it. The obtained crude extract further collected in amber colored screw cap bottle and stored in cooling condition till further use.

Detection and Characterization of Oil

The crude oil obtained through the extraction were further analyze by measuring gravity and density of the crude extract [14]. The obtained crude extract was further analyzed through thin layer chromatography [15], saponification [16], and estimation of oil content in it [17].

Evaluation of Microbial Contaminations


The obtained seed oil allowed to exposed microbial contaminants present in the laboratory. The obtained microbes were detected by microscopic method with the identification of morphological key markers [18].

Results and Discussion

The collected seeds were sundried divided into two equal halves. Half amount of seeds around 50 gm were soaked in distilled water for few days and remaining half amount of seeds were boiled in water for 3-4 hours. The extract obtained from soaked and boiled seeds appears to be brownish in color because of its seed coat color. The oil content present in the

seed sample were found to be floating on the water and later on separation extracted oil were done with the ether extraction using separating funnel. Seeds have medicinal values, and it also contains very good amount of natural oil in it [8]. The oil components were further has been confirmed with the help of thin layer chromatography.

The oil produced by the seeds were gradually accumulated at the upper surface of boiling water or normal water (Figure 1). The oil collected at the surface of boiling water were further mixed with petroleum ether and separated with separating funnel (Figure 2). The oil extract were further concentrated by the removal of petroleum ether by keeping the temperature of heating mantle at 45 °C around the boiling point of petroleum ether. The distillation helps to remove extra solvent present in the extract. The obtained extract were further characterized by measuring lipid content, specific gravity, density, protein, and glucose content. The compounds present in the ether extract were further analyzed and confirmed through the thin layer chromatography. The obtained extract is also subjected to the anti-microbial activity. In an anti-microbial assay, we found that instead of killing or inhibiting the growth of microbes, in the presence of extracted oil, confluent microbial growth were observed in comparison to control. There may be some nutrient component present in the extract responsible for enhanced microbial growth.



Eurasian Journal of
Science and Technology

Figure 1 Extraction of oil from plant seeds and further separation with separating funnel

Eurasian Journal of
Science and Technology**Figure 2** Determination of lipid content of oil by Saponification method

The detection of lipid content in the ether extract was done through the esterifications, during the esterification, we observed the formations of some unknown compounds and the probability of this compound could be Fatty Acid Methyl Esters (FAME) [19] (Figure 3). The estimation of lipid content in the extracted oil were found around 4.88 %, similarly in vegetable oil blend, other researchers found the oil content in vegetable oil blend around oleic acid (C18:1n-9; 63.3%), linoleic acid (C18:2n-6; 4.7%), and linolenic acid (C18:3n-6; 5.1%) [20]

Major components of the natural oils are terpenoids and phenolic compounds [21]. The natural and essential oil obtained from the plants or plants parts are basically composed of several chemical compounds or secondary metabolites such as terpenoids, shikimates, polyketides, and alkaloids which are produced through the shikimate pathways. These compounds have potential role in the food, cosmetic, and pharmaceutical industries (Figure 4) [6].


Eurasian Journal of
Science and Technology**Figure 3** Detection and separation of some unknown esterified product after esterification

The analysis of crude oil extract was done through the thin layer chromatography, in the chromatogram we observed uniform banding pattern in all samples of the crude oil (Figure 5). The samples seems to have pure oil components in the plant seeds with Rf values of

0.24, 0.45, 0.54, and 0.72. In the characterization of oil yielding seed like soybean, it was observed the oil components were observed at their respective Rf value such as Phospholipids at 0.01 ± 0.002 , Monoacylglycerides at 0.04 ± 0.013 ,


Triacylglycerides at 0.20 ± 0.034 , Tocopherols at 0.66 ± 0.105 , and Fatty acid esters at 0.95 ± 0.14 [22]. The characterization of secondary metabolites in the *Asplenium indicum* also have done through the thin layer chromatography [23]. In India, we have around sixteen edible oil yielding plants and numerous plants known for the extraction of medicinal oil. The oil can be

extracted from different plant parts such as root and shoot (leaves, fruits, seeds, bark, wood, flowers, and buds). The most of secondary metabolites are excluded from essential oils of the plant such as cellulose, glycerides, starches, sugars, tannins, salts, and minerals. The yield of the oil from the plant sources is ranges from 0.05-18.0%.



Eurasian Journal of
Science and Technology

Figure 4 Isolation and estimation of lipid content in oil extracts



Eurasian Journal of
Science and Technology

Figure 5 Analysis of crude oil extract by thin layer chromatography

The difference in the density of the crude oil extracts were observed in comparison to the water sample (Figure 6). The 5 ml sample shows the density of crude oil extract was 4.6 gm/cm^3 . The natural density of vegetable or plants oils comes in the range of $700\text{-}950 \text{ kg/m}^3$. Hence, during the oil extraction we find oil floating over the water. The freezing of such oils at lower temperature can indicate the presence of polyunsaturated fatty acids in it. Similar results we obtained during the experiments we kept soaking the seeds for the several days at low temperature, the

components present in the mixture are frozen over there. The density and viscosity of the sample plays an important role in the efficiency, lubrications, and functioning.

The crude oil obtained from seeds soaking in normal water labeled as sample 1, while sample 2 and sample 3 were obtained from boiling water. In one of the experiments, the sample viscosity was qualitatively done with the help of glass balls, we allowed to sink the glass ball through the test tube filled with sample. The ball present in sample takes more time in

comparison to water and acetone. We have three samples of extract sample 1 shows 2.5 sec, sample 2 shows 3.2 sec, sample 3 shows 5.1 sec, water shows 1.8 sec, and acetone shows 2.3 sec. Among this value, we can easily predict that the sample 3 of crude oil extract is more viscous in comparison water and acetone and other crude samples. The solvent viscosity is directly depending on the temperature, we found that

the natural oil obtained in this experiment having higher viscosity in comparison to water and alcohol. Viscosity is the important parameter for the process development of fatty acids in the industry or we can say it is important screening parameter for the elution of fatty acids from the columns. The viscosity of the liquid also helps in the designing heat transfer equipments.




Eurasian Journal of
Science and Technology

Figure 6 The difference in the density of the crude oil extracts in comparison to the water sample

During our investigation, we observed several contaminations of the microbes on our crude oil extracts [24]. Dominant contaminants are *Candida* and *Aspergillus* in our samples (Figure 7). The fungal contaminations such as yeast and mould is prevalent in vegetable edible oil [25], similar contamination is also observed in the case of oil yielding seeds [26]. The major prevalence of microbes was observed in stored

products of refinery in which significant were detected [27]. These sort of contaminations are not only observed in extracted oil, but also in dried leaves which are used for the tea bagging. The microbes having potential to produce lipase are the most prominent microbes that can be observed on oil obtained from plants or natural resources [28].



Eurasian Journal of
Science and Technology

Figure 7 Microscopic images of contaminated microbes in crude oil extracts

Conclusion

The plant of white Popinac or leucocephala has versatile characteristics, hence it has wide application in medicinal, cosmetics, pharmaceuticals, and confectionary to timber, almost all field. The plant is fast growing having higher photosynthetic rate; this plant can be an alternative to increasing deforestation and increasing global warming. Along rapid growth plant also provides timber for the furniture industry. From roots to shoots, each and every part of the plant has special use in human life such as food and medicine. The plant thought to be miracle timber tree as it is rapidly producing greater biomass; in folk medicine, it is to be used for the ruminants to increase the milk production and weight gain and have therapeutic value such as anthelmintic, antidiabetic, and also it is used to manage stomachache, contraception, and abortifacient. The major secondary metabolites present in this plant are alkaloid, cardiac glycosides, tannins, flavonoids, saponins, and Glycosides. In this investigation, we found that the plants taken into consideration are highly oil secreting with potential amount of nutrient content. The obtained oil has good amount of lipid content around 4.88% and fatty acids. In thin layer chromatography, we detected the presence of numerous fatty acids by comparing Rf values such as Phospholipids at 0.01 ± 0.002 , Monoacylglycerides at 0.04 ± 0.013 , Triacylglycerides at 0.20 ± 0.034 , Tocopherols at 0.66 ± 0.105 , and Fatty acid esters at 0.95 ± 0.14 . Also, we observed that the oil sample obtained from boiling water with higher viscosity in comparison to normal water sample.

Acknowledgements

The authors are gratefully acknowledged the School of Life Sciences, Swami Ramanand Teerth Marathwada University, Nanded 431606 for laboratory and other necessary facilities..

Consent for Publications

All authors read and approved the final manuscript for publication.

Availability of data and material

All the data are embedded in this article.

Funding

There was not any financial support for this study.

Conflict of Interest

The authors declare that they have no conflict of interest.

Ethics approval and consent to participate

No actual animal studies were performed in the present investigations.

ORCID

Jaya B. Puri

<https://www.orcid.org/0009-0008-2473-6676>

Abdulbasit Shaikh

<https://www.orcid.org/0000-0003-3280-5981>

Umesh Pravin Dhuldhaj

<https://www.orcid.org/0000-0002-7439-4669>

References

- [1] V.D. Samarth, S.R. Sinkar, U. Pandya, U.P. Dhuldhaj, Phytohormonal effect on in vitro callus induction of *Tinospora cordifolia* (Willd.) Miers, *Plant Cell Biotechnology and Molecular Biology*, 2019, 1383 [Google Scholar]
- [2] Swapnil P., Meena M., Singh S.K., Dhuldhaj U.P., Marwal A., Vital roles of carotenoids in plants and humans to deteriorate stress with its structure, biosynthesis, metabolic engineering and functional aspects. *Current Plant Biology*, 2021, 26:00203 [Crossref], [Google Scholar], [Publisher]
- [3] Dhawle K., Dhuldhaj U., Mulani R., Pharmacognostic studies and anatomical peculiarities in medicinal plant *Enicostemma axillare*. *Asian Journal of Ethnobiology*, 2021, 4:23 [Crossref], [Google Scholar], [Publisher]
- [4] Zayed M.Z., Sallam S.M.A., Shetta N.D., Review article on *Leucaena leucocephala* as one

- of the miracle timber trees. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2018, **10**:1 [Crossref], [Google Scholar], [Publisher]
- [5] Chowtivannakul P., Srichaikul B., Talubmook C., Antidiabetic and antioxidant activities of seed extract from *Leucaena leucocephala* (Lam.) de Wit. *Agriculture and Natural Resources*, 2016, **50**:357 [Crossref], [Google Scholar], [Publisher]
- [6] Chandran H., Meena M., Barupal T., Sharma K., Plant tissue culture as a perpetual source for production of industrially important bioactive compounds. *Biotechnology Reports*, 2020, **26**:e00450 [Crossref], [Google Scholar], [Publisher]
- [7] Rosida D.F., Djajati S., Nilamayu Z.A., Antibacterial Activity of *Leucaena leucocephala* Extracts on Growth of *Escherichia coli*. *Advanced Science Letters*, 2018, **23**:12268 [Crossref], [Google Scholar], [Publisher]
- [8] Nehdi I.A., Sbihi H., Tan C.P., Al-Resayes S.I., *Leucaena leucocephala* (Lam.) de Wit seed oil: Characterization and uses. *Industrial Crops and Products*, 2014, **52**:582 [Crossref], [Google Scholar], [Publisher]
- [9] Zayed M.Z., Samling B., Phytochemical constituents of the leaves of *Leucaena leucocephala* from Malaysia. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2016, **8**:174 [Crossref], [Google Scholar], [Publisher]
- [10] Sharma P., Kaur A., Batish D.R., Kaur S., Chauhan B.S., Critical insights into the ecological and invasive attributes of *Leucaena leucocephala*, a tropical agroforestry species. *Frontiers in Agronomy*, 2022, **4**: 890992 [Crossref], [Google Scholar], [Publisher]
- [11] Soetjipto H., Pradana R.C., Kristijanto A.I., April. Fatty acid composition and total lipid content of the seed oil of *Leucaena leucocephala* (Lam.) de Wit. In *IOP Conference Series: Materials Science and Engineering* IOP Publishing, 2019, **509**:012046 [Crossref], [Google Scholar], [Publisher]
- [12] Idowu S.O., Adeyemo M.A., Ogbonna U.I., Engineering and validation of a novel lipid thin film for biomembrane modeling in lipophilicity determination of drugs and xenobiotics. *Journal of Biological Engineering*, 2009, **3**:1 [Crossref], [Google Scholar], [Publisher]
- [13] Sundrasegaran S., Mah S.H., Extraction Methods of Virgin Coconut Oil and Palm-pressed Mesocarp Oil and their Phytonutrients. *EFood*, 2020, **1**:381 [Crossref], [Google Scholar], [Publisher]
- [14] Wang Z.M., Zhang J., Corrosion of multiphase flow pipelines: the impact of crude oil. *Corrosion Reviews*, 2016, **34**:17 [Crossref], [Google Scholar], [Publisher]
- [15] Mujeeb F., Khan A.F., Bajpai P., Pathak N., Phytochemical study of *Aegle marmelos*: Chromatographic elucidation of polyphenolics and assessment of antioxidant and cytotoxic potential. *Pharmacognosy Magazine*, 2017, **13**:S791 [Crossref], [Google Scholar], [PDF]
- [16] Patterson H.B.W., Hydrogenation of Fats and Oils (Second Edition) Theory and Practice, 2011, 329 [Crossref], [Google Scholar], [Publisher]
- [17] Wei F., Gao G.Z., Wang X.F., Dong X.Y., Li P.P., Hua W., Wang X., Wu X.M., Chen H., Quantitative determination of oil content in small quantity of oilseed rape by ultrasound-assisted extraction combined with gas chromatography. *Ultrasonics Sonochemistry*, 2008, **15**:938 [Crossref], [Google Scholar], [Publisher]
- [18] Senanayake I.C., Rathnayaka A.R., Marasinghe D.S., Calabon M.S., Gentekaki E., Lee H.B., Hurdeal V.G., Pem D., Dissanayake L.S., Wijesinghe S.N., Bundhun D., Morphological approaches in studying fungi: Collection, examination, isolation, sporulation and preservation. *Mycosphere*, 2020, **11**:2678 [Crossref], [Google Scholar], [PDF]

- [19] Widiarti N., Ni'mah Y.L., Bahruji H., Prasetyoko D., Development of CaO from natural calcite as a heterogeneous base catalyst in the formation of biodiesel. *Journal of Renewable Materials*, 2019, **7**:915 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [20] Guidoni M., de Christo Scherer M.M., Figueira M.M., Schmitt E.F.P., De Almeida L.C., Scherer R., Bogusz S., Fronza M., Fatty acid composition of vegetable oil blend and in vitro effects of pharmacotherapeutical skin care applications. *Brazilian Journal of Medical and Biological Research*, 2019, **52** [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [21] J.P. Fokou, P.M. Dongmo, F.F. Boyom, Int Open, 2020
- [22] Shafigullin D., Kask S., Gins M., Pronina E., Demyanova-Roy G., Soldatenko A., Fatty oil accumulation in vegetable soybean seeds and its thin-layer chromatography, *Agronomy Research*, 2021, **19**:586 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [23] Jadhav D., Ghatage M., Isolation and Characterization of Secondary Metabolites of *Asplenium indicum* by Using Thin Layer Chromatography. *Journal of Scientific Research*, 2021, **65** [[Crossref](#)], [[Google Scholar](#)], [[PDF](#)]
- [24] Ratho M.G., Dhotare J.M., Kamble G.T., Dhawale P.I., Dhuldhaj U.P., Chakke A.B., Puri J.B., Pathak A.P., Influence of various substrate treatment methods on yield and biological efficiency of *Pleurotus florida* mushroom. *World Journal of Advanced Engineering Technology and Sciences*, 2023, **8**:216 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [25] Okpokwasili G.C., Molokwu C.N., Yeast and mould contaminants of vegetable oils. *Bioresource Technology*, 1996, **57**:245 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [26] Bhat R., Reddy K.R.N., Challenges and issues concerning mycotoxins contamination in oil seeds and their edible oils: Updates from last decade. *Food Chemistry*, 2017, **215**:425 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [27] Gaylarde C.C., Bento F.M., Kelley J., Microbial contamination of stored hydrocarbon fuels and its control. *Revista de Microbiologia*, 1999, **30**:1 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [28] Popoola B.M., Onilude A.A., Microorganisms associated with vegetable oil polluted soil. *Advances in microbiology*, 2017, **7**:377 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]